

CrossRef DOI of original article:

Laceration of the Left Renal Pelvis in a Hydronephrotic Kidney After Blunt Abdominal Trauma -A Case Report

Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970

Abstract

Index terms—

1 I. INTRODUCTION

Injuries caused by traffic accidents are responsible for approximately 1.35 million of deaths worldwide and 50 million of non-fatal injuries, most of which the people involved ends up being incapacitated. In Brazil, trauma is the third leading cause of overall mortality, with injuries occurring in traffic accounting for 40,000 deaths and more than 150,000 victims being injured with a high degree of severity annually. The incidence is more prevalent in males and young adults, mainly between 15 and 44 years of age. 7,8 The kidney is the organ of the genitourinary tract most frequently injured during trauma, which affects up to 10% of patients hospitalized for abdominal trauma. Renal trauma mainly affects men, in 72-93% of cases, it is more frequent in the younger population, between 31 and 38 years old, and pathological polycystic kidneys, with tumors or with hydronephrosis secondary to stenosis of the ureteropelvic junction are more prone to trauma, even in lower energy impacts. 1,2 London Journal of Medical and Health Research Kidney trauma can occur through three mechanisms: blunt trauma, penetrating trauma and high-speed deceleration. Blunt trauma is responsible for 71-95% of kidney injuries, the cause of which comes mainly from car accidents and falls. Most kidney injuries are mild, while severe trauma is more prevalent in patients with penetrating trauma than in those who have suffered blunt trauma (27-68% vs. 4-25%). 1,2 The most used classification of renal trauma is the one proposed by the American Association for the Surgery of Trauma (AAST), revised in 2011, which grades renal injuries from I to V with progressive severity (Table 1).

2 GRADE II

Parenchymal laceration < 1 cm in depth, without urinary extravasation, and with non-expanding perirenal hematoma.

3 GRADE III

Laceration > 1 cm deep, without collecting system damage and without urinary extravasation.

4 GRADE IV

Laceration in the cortex, medulla and collecting system, possibly with renal artery or vein injury, with contained hemorrhage.

5 GRADE V

Kidney is completely fragmented (shattered) and there is avulsion of the renal hilum, which cuts off the organ's blood supply.

Source: The patient was brought in on a rigid board, with a cervical collar, his airways were patent and there was no pain on palpation of the cervical spine. Upon examination of the chest, expansion was preserved, there was no crackling or pain on palpation, and breath sounds were present bilaterally. Flaccid abdomen, diffusely painful on palpation, without signs of peritoneal irritation, with present bowel sounds and stable pelvis. The

7 III. DISCUSSION

41 pupils were isochoric and isophotoreactive and the Glasgow Coma Scale was 15. At the time of hospital care,
42 the patient was conscious and oriented in time and space, without neurological deficits, hemodynamically stable,
43 reporting left abdominal pain. He denied traumatic brain injury, headache, nausea, vomiting and amnesia.

44 The victim denied comorbidities and use of continuous medication and was unaware of previous kidney
45 pathology. He also denied tobacco use but reported moderate alcohol consumption (6 cans per weekend). There
46 was no family history of neoplasms or organ malformations.

47 Computed tomography (CT) scans of the skull, chest and abdomen were requested. According to the high-
48 energy trauma protocol established by the service, tomography is available 24 hours a day and is readily accessible,
49 in addition, the patient was hemodynamically stable and needed imaging tests to complement the evaluation,
50 opting for CT instead of FAST (Focused Assessment with Sonography for Trauma).

51 The computed tomography report of the abdomen (Figure 1) showed a large retroperitoneal collection adjacent
52 to the left kidney, showing pelvis distention and the "claw sign". He also considered that the possibility of a
53 large adjacent urinoma could be a differential diagnosis, suggesting further investigation. The spleen had normal
54 dimensions, smooth contours and homogeneous attenuation. The exam also showed a small ascites, and there were
55 no other significant alterations, including in the liver, bile ducts, pancreas, adrenals, retroperitoneal lymph nodes
56 or in retroperitoneal vessels, with those cited being found in anatomical aspect and normal dimensions. Chest
57 and skull tomography showed no alterations. From the data found in the anamnesis, physical examination and
58 CT, it was concluded that it was a renal lesion grade V. It wasn't visualized ureteral lesion in the imaging exams
59 due to massive leakage of urine and total distortion of the renal anatomy. Therefore, the patient was referred and
60 underwent exploratory laparotomy, left nephrectomy and distal ureteral ligation. During the surgical procedure,
61 no intraperitoneal free fluid or changes in hollow viscera were found.

62 6 London Journal of Medical and Health Research

63 However, when exploring the left retroperitoneal region, a large amount of urine was drained and a hyperdistended
64 renal capsule was found, in addition to the kidney with a grade V lesion, characterized by complete laceration
65 with avulsion of the hilum and devascularization. In addition, the ureter was patent in the renal hilum, with a
66 complete section 10 cm from the ureteropelvic junction, with a grade V urethral lesion, and a distal ligation was
67 performed. No other injuries were found. The anatomical specimen removed from the patient during surgery is
68 shown in The product of the nephrectomy was sent for anatomopathological study, whose report showed signs of
69 rupture of the renal tissue, with ischemic changes and hydronephrosis, in addition to the absence of malignancy,
70 while in the ureter, focal stenosis and complete section of this were observed, suggesting stenosis of the UPJ
71 (junction pyelic ureter).

72 7 III. DISCUSSION

73 In the case observed, the patient's kidney was previously hydronephrotic due to UPJ stenosis, despite not having
74 a previous diagnosis, a fact that increases the risk of more serious injuries and the need for nephrectomy in blunt
75 trauma. 3 However, despite the spleen being the most affected organ in this type of trauma, the patient did
76 not have splenic injury. 4 However, there was ureter trauma, which is extremely uncommon in blunt trauma,
77 whereas ureter injury is usually secondary to iatrogenesis in surgical procedures or due to penetrating trauma,
78 being the most common blunt injury in children with congenital anomalies of the urinary tract, such as UPJ
79 stenosis, which causes hydronephrosis. Ureteral trauma, when identified during surgery, should be treated as
80 early as possible by surgical correction. 9 Kidney trauma should be suspected whenever there is a history of an
81 event with fast deceleration or direct impact on the flank region, back, lower chest and upper abdomen. In these
82 cases, the physical examination may reveal bruises on the flank or upper abdomen, signs of peritoneal irritation,
83 a palpable mass, ecchymosis or abrasions, and rib fractures. In this context, CT with intravenous contrast is
84 the imaging method of choice for hemodynamically stable patients who are victims of both penetrating trauma
85 and blunt trauma, as it allows the definition of the location and severity of the injury, as well as allows the
86 visualization of associated traumas.

87 FAST, on the other hand, is useful to demonstrate the presence of free fluid in the cavity, but it is inferior to CT
88 in terms of resolution and ability to define renal trauma. 1,6 The management of the patient victim of renal trauma
89 has as priority the control of blood loss, the preservation of the renal tissue and the prevention of complications.
90 While in the past it was thought that the best way to control bleeding London Journal of Medical and Health
91 Research and prevent nephrectomies was through surgery, in recent decades the management of renal trauma has
92 evolved towards prioritizing nonoperative approaches. 1 A recent meta-analysis has shown that the management
93 of non-operative treatment (NOT) is the treatment of choice for low-grade renal trauma, and it should also
94 be considered the first-line treatment of high-grade blunt trauma and penetrating trauma, as it is associated
95 with lower mortality rates, shorter hospital stay hospitalization and similar morbidity rates. 4,5 Absolute
96 indications for surgery involve: hemodynamic instability and unresponsiveness to aggressive resuscitation due
97 to renal hemorrhage, grade V vascular injury, and expanding or pulsating perirenal hematoma found during
98 laparotomy performed due to associated injuries to other organs. Relative indications for surgery include, among
99 others: large urine extravasation (spontaneous resolution in most cases), presence of non-viable renal tissue and
100 arterial thrombosis installed for more than 4 hours. 1,2

8 IV. CONCLUSION

It was possible to conclude that patients with previous renal alterations are more susceptible to more severe injuries, even in blunt trauma, where the most affected organ would be the spleen.

Injury to the ureter is extremely uncommon in blunt trauma, the main cause of which is iatrogenic. The case described above is atypical because the patient had left kidney injury, which is anatomically close to the spleen, but did not present injury to this organ, but a complete section of the ureter. For the definitive diagnosis, the correlation between data from the anamnesis, physical examination, imaging tests, intraoperative findings and the result of the anatomopathological analysis was essential.

9 Attachments

10 GRADE II

Parenchymal laceration < 1 cm in depth, without urinary extravasation, and with non-expanding perirenal hematoma.

11 GRADE III

Laceration > 1 cm deep, without collecting system damage and without urinary extravasation.

12 GRADE IV

Laceration in the cortex, medulla and collecting system, possibly with renal artery or vein injury, with contained hemorrhage.

13 GRADE V

Kidney is completely fragmented (shattered) and there is avulsion of the renal hilum, which cuts off the organ's blood supply.

Source: ^{1 2}



Figure 1: Figure 1 :

121

¹ Laceration of the Left Renal Pelvis in a Hydronephrotic Kidney After Blunt Abdominal Trauma -A Case Report © 2023 Great | Britain Journals Press || Volume 23 Issue 1 ??? Compilation 1.0

² Laceration of the Left Renal Pelvis in a Hydronephrotic Kidney After Blunt Abdominal Trauma -A Case Report || © 2023 Great | Britain Journals Press Volume 23 Issue 1 ??? Compilation 1.0

2

Figure 2: Figure 2 .



2

Figure 3: Figure 2 :



1

Figure 4: Figure 1 :

ABSTRACT

Introduction: The kidney is one of the most frequently injured organs during blunt abdominal trauma, and the probability increases significantly if there is any renal pathology, such as hydronephrosis. The ureter Injury, which is extremely uncommon in blunt trauma, also increases its incidence in the presence of previous renal pathologies. Case report: Patient J7R.A, male, 30 years old, brought to the HMMSJP hemodynamically stable by SIATE after suffering a automobile x truck accident as a driver and the

hemodynamically unstable.
Conclusion:
Blunt kidney injury is more common in patients with previous renal alterations, while ureteral injuries are mostly due to iatrogenic events during surgery or penetrating trauma, being uncommon in blunt trauma. The correlation between anamnesis, physical examination and imaging tests is essential for the appropriate and individualized management of the patient.

1

Graduation of Kidney Injury	Description of Injury
GRADE I	Subcapsular hematoma, non-expanding, without laceration in the parenchyma, and there may be microscopic or macroscopic hematuria, but without changes in urine tests.

3 London Journal of Medical and Health Research 2 Laceration of the Left Renal Pelvis in a Hydronephrotic Kidney After Blunt Abdominal Trauma -A Case Report

Figure 6: Table 1 :

2

Graduation of Ureteral Injury	Description of the Injury
GRADE I	Bruise; contusion or hematoma without devascularization.
GRADE II	Laceration; < 50% transection
GRADE III	Laceration; > 50% transection
GRADE IV	Laceration; complete transection with devascularization < 2cm
GRADE V	Laceration; avulsion with > 2cm of devascularization

Source: table adapted from the book "Fundamental Urology" by SBU, 2010
II. CASE REPORT

J. R. A., 30 years old, male, 1.85 m, 92 kg, is brought by SIATE after suffering an accident with a car x truck as a driver, with the car overturning.

Figure 7: Table 2 :

1

Graduation of Kidney Injury	Description of Injury
GRADE I	Subcapsular hematoma, non-expanding, without laceration in the parenchyma, and there may be microscopic or macroscopic hematuria, but without changes in urine tests.

Figure 8: Table 1 :

Figure 9:

2

Graduation of Ureteral Injury	Description of the Injury
GRADE I	Bruise; contusion or hematoma without devascularization
GRADE II	Laceration; < 50% transection
GRADE III	Laceration; > 50% transection
GRADE IV	Laceration; complete transection with devascularization < 2cm
GRADE V	Laceration; avulsion with > 2cm of devascularization

Source: table adapted from the book "Fundamental Urology" by SBU, 2010 London Journal of Medical and Health Research 7 Laceration of the Left Renal Pelvis in a Hydronephrotic Kidney After Blunt Abdominal Trauma -A Case Report © 2023 Great Britain Journals Press | | Volume 23 Issue 1 ??? Compilation 1.0

Figure 10: Table 2 :

-
- 122 [London Journal of Medical and Health Research] , *London Journal of Medical and Health Research*
123 [Filho ()] , M Z Filho . *Urologia. Editora Planmark* 2010. São Paulo.
- 124 [Silva ()] ‘Abordagem do trauma renal -artigo de revisão’. L F Silva . *Revista Do Colégio Brasileiro de Cirurgias*
125 2009. 36.
- 126 [Bjurlin ()] ‘Comparison of nonoperative and surgical management of renal trauma’. M A Bjurlin . *Journal of*
127 *Trauma and Acute Care Surgery* 2017. 82.
- 128 [Mingoli ()] ‘Operative and nonoperative management for renal trauma: comparison of outcomes’. A Mingoli .
129 *Therapeutics and Clinical Risk Management* 2017. 13.
- 130 [Erlich and Kitrey ()] ‘Renal trauma: the current best practice’. T Erlich , N D Kitrey . *Therapeutic Advances*
131 *in Urology*, 2018. 10.
- 132 [Serafetinides ()] ‘Review of the Current Management of Upper Urinary Tract Injuries by the EAU Trauma
133 Guidelines Panel’. E Serafetinides . *European Urology* 2015. 67.
- 134 [Buckley and Mcaninch ()] ‘Revision of current American Association for the Surgery of Trauma renal injury
135 grading system’. J C Buckley , J W Mcaninch . *Journal of Trauma* 2011.
- 136 [Malta ()] ‘Road traffic injuries and deaths and the achievement of UN Sustainable Development Goals in Brazil:
137 results from the Global Burden of Disease Study’. D C Malta . *Revista da Sociedade Brasileira de Medicina*
138 *Tropical* 1990 to 2019. 2022. 55.
- 139 [Ibiapino ()] *Serviço de atendimento móvel de urgência: epidemiologia do trauma no atendimento pré-hospitalar.*
140 *Revista da Faculdade de Ciências Médicas de Sorocaba*, M K Ibiapino . <[https://revistas.pucsp.br/
141 index.php/RFCMS/article/view/30805/pdf](https://revistas.pucsp.br/index.php/RFCMS/article/view/30805/pdf)> 2017.